Lattice study of
mesons near deconfining transition

Hideo Matsufuru (Hiroshima Univ.)

for QCD-TARO Collaboration

23 March 1999
Workshop on Hadron Physics
Hiroshima Univ. (Higashi-Hiroshima)

— *CONTENTS* —

Introduction
Our Approach and Strategy
Lattice Setup
Results of Simulation:
    (1) t-correlators
    (2) Wave functions
Summary and Outlook

The QCD-TARO Collaboration

Ph. de Forcrand[a], *M. García Pérez*[b], *T. Hashimoto*[c], *S. Hioki*[d], *H. Matsufuru*[e], *O. Miyamura*[e], *A. Nakamura*[f], *I.-O. Stamatescu*[g,h], *T. Takaishi*[i] *and T. Umeda*[e]

[a] SCSC, ETH-Zürich, Switzerland
[b] Dept. Física Teórica, Univ. Autónoma de Madrid, Spain
[c] Dept. Appl. Phys., Fac. Engineering, Fukui Univ., Japan
[d] Dept. of Physics, Tezukayama Univ., Nara, Japan
[e] Dept. of Physics, Hiroshima Univ., Japan
[f] Res. Inst. for Inform. Sci. and Education, Hiroshima Univ., Japan
[g] Inst. Theor. Physik, Univ. of Heidelberg, Germany
[h] FEST, Heidelberg, Germany
[i] Hiroshima University of Economics, Japan

Physical goals:

- Finite Temperature QCD
- Finite Chemical Potential
- MCRG and Improved Actions

Introduction

Finite temperature hadron properties:

Change of masses and width near T_c
What happens on hadrons or what new effect above T_c ?

Example of effective theory:
NJL model analysis

c.f. Hatsuda and Kunihiro, Phys. Rep. 247 (1994) 221

- Mean-field theory
- $N_f = 2 + 1$
- Abobe T_c : Soft modes

Figure

Lattice QCD

Euclidean 4-D lattice

- Gauge field: link var. $U_\mu(x) \sim e^{-igA_{\mu(x)}}$
- Quark field: $\bar{\psi}(x)$, $\psi(x)$

$$\langle O \rangle = \int \mathcal{D}U\mathcal{D}\bar{\psi}\mathcal{D}\psi O[U,\bar{\psi},\psi]\exp(-S[U,\bar{\psi},\psi])$$

$$\Downarrow$$

Monte Carlo integration

At finite temperature:

- $N_s \gg N_t$, $T = 1/N_t a$
- Boundary condition
  - boson: periodic
  - fermion: anti-periodic

Lattice Results

At $T > T_c$:

- Spatial correlators ($\leftrightarrow$ Screening masses)
  Correlation in π, σ sectors is large ?
  Weak in other sectors ($\sim$ free quarks)
  Born et al. (1991) ($N_f = 4$)
  No bosonic pole
  Gupta (1992) (quenched: $N_f = 0$)

- Baryon number susceptibility
  Large, $\chi_S \sim \chi_{NS}$ (isosinglet and nonsinglet)
  $\Rightarrow$ Fundamental excitations are quarks ?
  Gottlieb et al. (1987) ($N_f = 2$)

- Temporal correlators ($\leftrightarrow$ Pole masses)
  - Temperature Green function
    $\Leftarrow$ correlator in Euclidean time direction
  - On anisotropic lattice
    Hashimoto, Nakamura and Stamatescu (1993)

Our Approach

From correlators in Euclidean time direction;

- Pole masses (compared with screening masses)
- Wave function
- Spectral function

Need detailed information in t- (temperature) direction

$\Rightarrow$ *Anisotropic lattice*

[Karsch (1982),
Burgers, Karsch, Nakamura and Stamatescu (1988)]

This work:

- coarse lattice, ($a_s^{-1} = 0.85$ GeV)
- unimproved actions

$\longrightarrow$ Qualitative result, Development of procedures

Strategy

Difficulty:

Mass is extracted at $t \gg 1$.

However, at $T > 0$, temporal extent is short.

$\Rightarrow$ Choice of hadronic operator is significant.

Investigate following questions.

(1) Define the "hadronic operator" as one which has sufficiently large overlap with corresponding states. Then, what happens on this operator at $T > 0$?

(2) Is there bound state at $T > T_c$?
Haw can we show that ?

(3) Develop reliable proceedure to extract the pole masses with short extent in t-direction.

Simulation Parameters

Lattice:

$12^3 \times N_t$,$\beta = 5.68$, $\gamma = 4.0$, quenched

$N_t = 72$ ($T \simeq 0$), 20 ($T < T_c$), 16, 12 ($T > T_c$) : $T = 1/N_t a_t$

- #conf. $= 60$
- Anisotropy: $\xi \equiv a_s/a_t = 5.3(1)$
  from the ratio of Wilson loops
  Engels, Karsch and Scheideler (1997), Klassen (1998)
- Cutoff: $a_s^{-1} = 0.85$ GeV, $a_t^{-1} = 4.5(2)$ GeV
  from heavy quark potential

Quark: Anisotropic Wilson action

- Hopping parameter and bare anisotropy:

κ_s	γ_F	m_q	m_{PS}	m_V $[GeV]$
0.0810	4.05	0.17	0.81	0.90
0.0840	3.89	0.12	0.68	0.80
0.0860	3.78	0.10	0.61	0.75

- γ_F determined by calibration
- Periodic b. c. for spatial direction

Correlators

Measure the wave function

$$w_\Gamma(r,t) = \sum_{\vec{x}} \langle \bar{q}(\vec{x}+\vec{r},t)\Gamma q(\vec{x},t) O_\Gamma^\dagger \rangle$$

where,

$$O_\Gamma(y) = \bar{q}^{(\varphi')}(y)\Gamma q^{(\varphi)}(y),$$

$$q^{(\varphi)}(y) = \sum_{\vec{r}} \varphi(\vec{r}) q(\vec{y}+\vec{r}, y_4)$$

Γ: γ_5 (Ps), γ_1 (V), 1 (S), $\gamma_1\gamma_5$ (A)
$\varphi(\vec{r})$: smearing function

We use "exp" smearing function:

$$\varphi(\vec{r}) = \exp(-ar^p)$$

with a, p extracted from observed wave function as well as "point" source: $\varphi(\vec{r}) = \delta_{\vec{r},0}$.

$\Rightarrow$ three sets of source smearing:
"point-point", "point-exp", "exp-exp"

Smearing Function

"exp" smearing function:

$$\varphi(\vec{r}) = \exp(-ar^p)$$

$\rightarrow$ "exp-exp" source smearing corresponds to the convolution of two $\varphi(r)$

$$\varphi^{(2)}(\vec{r}) = \int d^3x \varphi(\vec{r}+\vec{x})\varphi(\vec{r})$$

$$\kappa_s = 0.081 \ (m_q \sim 2m_s)$$



t-correlators

Correlators in Euclidean time direction:

$$C_\Gamma(t) = w(\vec{r}=0,t) = \sum_{\vec{x}} \langle O_\Gamma(\vec{x},t) O_\Gamma^{(\varphi)\dagger}(0,0)\rangle$$

Effective mass: $m^{(eff)}$ s.t.

$$\frac{C(t)}{C(t+1)} = \frac{\cosh[m^{(eff)}(N_t/2-t)]}{\cosh[m^{(eff)}(N_t/2-t-1)]}$$

For large t, $m^{(eff)} \to m_\Gamma$

Problem at finite temperature:

- Temporal extent is not enough for $m^{(eff)} \to m_\Gamma$
- Even in free quark case, ficticious plateau is observed

Effective Mass Plots at $N_t = 72$

$$\kappa_s = 0.081 \ (m_q \sim 0.17 GeV)$$

Pseudoscalar



Vector



Effective Mass Plots at $T > 0$

At $N_t = 72$, "exp-exp" correlators almost have required property: large overlap with meson states.

In the following analysis, "exp-exp" correlators are used.

$\kappa_s = 0.086\ (m_q \sim 0.10 GeV)$



Large qualitative change at $T > T_c$

Chiral symmetry restoration (?)

Answer to (1)

At $N_t = 72$, "exp-exp" correlator almost have
required property: large overlap with meson states
(More detailed analysis — e.g. Variational analysis)

$\Downarrow$

Observe the temperature dependence of them:
"masses" extracted from exp-exp correlators
(assuming the existence of bound states even above T_c)

At finite temperature,
- correlators with various sources do not give the same effective mass
- Difficult to find clear plateau

Uncertainty of extracted "mass" values are large (20–30 %)

Temperature dependence of "masses"

$\kappa_s = 0.081\ (m_q \sim 0.17GeV)$



$\kappa = \kappa_c\ (m_q \to 0 \text{ limit})$



t-dependence of the wave function

$$w_\Gamma(r,t) = \sum_{\vec{x}} \langle \bar{q}(\vec{x}+\vec{r},t)\Gamma q(\vec{x},t) O^\dagger(0)\rangle$$

If there is no bound state (like free quark case), wave function become broader as t.

In this case, $\phi_\Gamma(r,t) = w_\Gamma(r,t)/w_\Gamma(r=0,t)$

(normalized at spatial origin)

increase as t.



Answer to (2)

The wave function with "exp-exp" source become narower as t at all T.

- $\phi(r,t)$ should approaches certain shape
- exp-exp source (convolution of two φ(r)) is too broad

It sugests there are bound states even above T_c.

Further analysis:
- Wave functions with various sources
- Mix point-exp and exp-exp correlators to give flat t-dependence

Outlook on (3)

Possible analysis:
◦ Variational analysis
◦ using the wave function
— *in progress*

Spectral function may have broad width at $T > 0$
In this case, extraction of mass as the peak position of spectral function is difficult

⇓

Direct determination of the spectoral function from correlator:
— *Study in progress*
Nucl.Phys.B(Proc.Suppl.)63(1998)460 (Lattice 97).

Summary

(1) We observed the temperature dependence of "exp-exp" correlator, as the best operator at $T = 0$. Clear T-dependece of exracted masses were observed. Uncertainty of them are rather large at $T > 0$. ☺

(2) Observed t-dependence of the wave function suggest there is certain bound state up to 1.5 T_c. Further verification is necessary. ☺

(3) Now under progress. For example, direct determination of the spectral function from the correlator is investigated.

Outlook

Other channels: baryons etc.
Heavy quarkonium: J/ψ, ...
Relation with topological quantities
Precise determination of mass shift
Spectral function

$\Rightarrow$ Larger, improved lattice

With dynamical quarks