Spectral function with maximum entropy method on anisotropic lattice

Takashi Umeda, (CCP, Univ. Tsukuba)
Hideo Matsufuru, (YITP, Kyoto Univ.)

JPS meeting
22–25 September 2001, Okinawa Int. Univ.

Introduction

Hadron properties at Finite Temperature:

What happens on hadrons

or what new effects above T_c ?

Quark Gluon Plasma formation

$\Downarrow$

- J/ψ suppression above T_c

  Matsui and Satz (1986), Hashimoto et al (1986)

- Mass shift near T_c

  Hashimoto et al (1986)

Spectral Function at $T > 0$

from Maximum Entropy Method

Contents

1. Maximum Entropy Method
   - 1-i: Outline of MEM
   - 1-ii: Singular Value Decomposition

2. Approach for $T > 0$

3. Results
   - 3-i: Simulation parameters
   - 3-ii: Spectral function at $T = 0$
   - 3-iii: nt_{max} dependence
   - 3-iv: Spectral function at $T > 0$

4. Discussion

Maximum Entropy Method (MEM)

Reconstruction of a Spectral Function with MEM

Asakawa et al. (1999)

$$C(\tau) = \int d\omega K(\tau, \omega) A(\omega)$$

kernel : $K(\tau, \omega) = e^{-\omega\tau} + e^{-\omega(T-\tau)}$

standard χ^2-fit $\Longrightarrow$ ill-posed problem

$\Longrightarrow$ MEM (based on Bayes' theorem)

$$\boxed{\text{Maximization of } Q = \alpha S - L}$$

$$S = \int d\omega \left(A(\omega) - m(\omega) - A(\omega) \ln \frac{A(\omega)}{m(\omega)}\right)$$

$m(\omega) = m_0 \omega^2$: default model

α : constant $\Rightarrow$ to be integrated out

L : Likelihood function (χ^2-term)

Singular Value Decomposition (SVD)

$$\begin{aligned} K(\tau,\omega) &= e^{-\omega\tau} + e^{-\omega(T-\tau)} \\ &= V(\tau,\tau')w(\tau',\tau'')U(\omega,\tau'')^t \end{aligned}$$

$w(\tau,\tau')$: diagonal matrix

$$A(\omega) = m_0\omega^2 \prod_{i=1}^{N} \exp\{b_i u_i(\omega)\}$$

$u_i(\omega)$ $(= U(\omega,\tau_i))$: basis in singular space

b_i : parameters

▷ Figure

- need for large N (fine temporal resolution)
- difficult to reconstruct

  narrow peak at $\omega \gg 1$

  wide peak at $\omega \ll 1$

Singular Value Decomposition (2)

$$A(\omega) = m_0\omega^2 \prod_{i=1}^{N} \exp\{b_i u_i(\omega)\}$$

$\exp\{u_i(\omega)\}$



Approach for Finite temperature(1)

- Large data points for reconstruction of $A(\omega)$

  $\Longrightarrow$ Anisotropic lattice

- Restriction from SVD
  - High statistics
  - Shift of zero point $\omega \longrightarrow \omega + dm$

$$e^{dm \cdot \tau} C(\tau) = \int d\omega \; K(\tau, \omega) A'(\omega)$$

$$\Longrightarrow A(\omega) = A'(\omega - dm)$$

(real*8 code works on heavy quarks)

- Check of nt_{max} dependence at $T = 0$

  τ-range : $1 \sim nt_{max}$

Approach for Finite temperature(2)

- Estimation for width of Spectral function



Simulation parameters

Gauge parameters

Anisotropic plaquette action $(a_\sigma/a_\tau = 4)$

Scale is determined from Sommer scale r_0

β	γ_G	size	a_σ^{-1} (GeV)
6.10	3.2108	$20^3 \times N_t$	2.030(13)

N_t	T/T_c	conf.#
160	~ 0	500
32	~ 0.9	500
26	~ 1.1	500

Quark parameters

$O(a)$ improved Wilson quark action

Clover coefficients : Tadpole improved tree level

	κ	γ_F	quark mass
κ_{light}	0.1235	4.034	$\sim$ strange quark
κ_{heavy}	0.1120	4.000	$\sim$ charm quark

Spectral function at T = 0



These are consistent with

pole fit or previous works

nt_{max} dependence $(T = 0)$



▷ MEM work $(nt_{max} \cdot a_\tau) \gtrsim 1fm$

It is difficult to reconstruct

from correlator with multi states

Spectral function at T > 0



Spectral function at T > 0



Not correctly reconstructed at T > 0

Discussion

It is difficult to reconstruct
a Spectral Function at $T > 0$ with MEM

- ▷ temporal resolution is enough
  $a_\tau \simeq 8\mathrm{GeV} \Longrightarrow 1/T_c \sim 28 a_\tau \quad (N \leq 14)$
- ▷ larger statics is needed
  present stat.$=$ 500conf. $\Longrightarrow$??
- ▷ MEM works $(nt_{max} \cdot a_\tau) \gtrsim 1\mathrm{fm}$

  $\Longrightarrow$ using smeared operators (?)
- ▷ Improvement of default model $m(\omega)$