Accurate determination of gauge and quark actions on anisotropic lattices

Hideo Matsufuru

High Energy Accelerator Research Organization (KEK)

Hidenori Fukaya, Tetsuya Onogi, Takashi Umeda

Yukawa Institute for Theoretical Physics, Kyoto Univ.

Masanori Okawa

Dept. Physics, Hiroshima Univ.

Web-site: http://www.rcnp.osaka-u.ac.jp/~matufuru/

JPS meeting autumn 2004

30 September 2004, Kochi

Introduction

Anisotropic lattice: $a_\tau < a_\sigma \Rightarrow$ Anisotropy $\xi = a_\sigma / a_\tau$

Useful in various subjects:

- Heavy quark physics (our main purpose)
  a relativistic framework which enables
  - continuum limit
  - systematic improvement
  - modest size of computation
- Finite temperature physics
  — better resolution in Euclidean temporal direction
  $\rightarrow$ larger number of Matsubara frequency
- Correlators which rapidly grows statistical noises
  glueballs, exotics, excited states, etc.
  — easy to follow changes in t

Introduction

Disadvantage:

- Needs nonperturbative calibration (tuning of parameters)
- Systematic errors due to anisotropy must be controlled.

Present statistial accuracy: O(2%) in parameters

$\rightarrow$ O(10%) accuracy in spectrum and matrix elements

— not sufficient for precision computation

Our goal: 0.2% level of accuracy for parameters

$\Rightarrow$ a few percent calculation of matrix elements

Present stage: in quenched approximation

— studies of calibration procedure

Target anisotropy: $\xi = 4$, $\beta = 5.7 - 6.3$ $(a_s^{-1} = 1 - 3$ GeV)

Calibration of gauge field

Previous work by Klassen: 1% level of statistical error.

T.R.Klassen, Nucl. Phys. B 533 (1998) 557.

— For present purpose, this accuracy is not sufficient.

Renormalized anisotropy $\xi_G \leftarrow$ static potential
We define ξ_G through hadronic radius r_0

r_0: $r_0^2 F(r_0) = 1.65$ ($F(r)$: force)

Sommer, Nucl. Phys. B411 (1994) 839

- Precise computation possible
  — Lüscher-Weisz noise reduction technique

  Lüscher and Weisz, JHEP 0109 (2001) 010.
- Good scaling behavior (if lattice scale is set by r_0)
- Continuum limit in terms of scale set by r_0
  $\rightarrow$ systematic error in γ_G disappears as $a \rightarrow 0$

Calibration of gauge field

Result at $\beta = 5.75$





$\gamma^* = 3.1406(55)$ (0.2% accuracy)

Calibration of gauge field

Result at $\beta = 6.0$





$\gamma^* = 3.2012(66)$ (0.2% accuracy)

Calibration of gauge field

Results (preliminary)



More accuracy is needed for high β region (in progress).

Global fit $\rightarrow$ γ_G^* with 0.2% level uncertainty

Estimate of systematic errors

Calibration of quark field

Quark action: $O(a)$ improved Wilson action

$$S_F = \sum_{x,y} \bar{\psi}(x) K(x,y) \psi(y)$$

$$\begin{aligned} K(x,y) \quad = \quad & \delta_{x,y} - \kappa_\tau \left[(1-\gamma_4) U_4(x) \delta_{x+\hat{4},y} + (1+\gamma_4) U_4^\dagger(x-\hat{4}) \delta_{x-\hat{4},y} \right] \\ & - \kappa_\sigma \sum_i \left[(r-\gamma_i) U_i(x) \delta_{x+\hat{i},y} + (r+\gamma_i) U_i^\dagger(x-\hat{i}) \delta_{x-\hat{i},y} \right] \\ & - \kappa_\sigma c_E \sum_i \sigma_{4i} F_{4i}(x) \delta_{x,y} + r \kappa_\sigma c_B \sum_{i>j} \sigma_{ij} F_{ij}(x) \delta_{x,y} \end{aligned}$$

$\gamma_F = \kappa_\tau / \kappa_\sigma$: bare anisotropy

γ_F, c_E and c_B: to be tuned nonperturbatively

Calibration of quark field

γ_F, c_E, c_B, κ_c, and c_A (for axial current) in massless limit.
— γ_F must be tuned most precisely (to $O(0.2\%)$ level)

Applied techniques:

- Nonperturbative renormalization technique

  Lüscher et al., Nucl. Phys. B478 (1996) 365, B491 (1997) 323

- Spectroscopy in fine and coarse directions

(1) Schrödinger functional method $\Rightarrow$ tuning of c_E, c_B

(2) Spectrum in coarse/fine directions on lattices of T,$L \sim$ 2 fm
Physical isotropy conditions for m_{PS}, m_V $\Rightarrow$ γ_F, (c_B)

(3) Schrödinger functional method $\Rightarrow$ c_A, κ_c

(4) Check of systematic errors
Light hadron spectrum and dispersion relation
Taking the continuum limit

Calibration of quark field

Result at $\beta = 6.10$
$8^3 \times 64$, $\xi = 4$
(γ_G by Klassen)
$a_s^{-1} = 2.0$ GeV



Quark mass defined by PCAC should not depend on kinematical parameters up to $O(a^2) \rightarrow$ improvement of c_E

- NP improved c_E is larger than tadpole tree value, $c_E = 1.26$.
- ΔM is not sensitive to c_B

Calibration of quark field

Determination of γ_F:



$\beta = 5.75$, $12^2 \times 24 \times 96$, $a_s^{-1} = 1.1$ GeV, 136 confs.

γ_F is determined from meson masses in fine/coarse directions

- precision of γ_F is still not sufficient
- consistent with γ_F from dispersion relation

Summary and outlook

We are developing calibration procedures of anisotropic lattices for precise computation of heavy-light matrix elements.

In quenched approximation,

- gauge field is calibrated with sufficient precision.
- quark field is now under investigation.
  - accurate determination of γ_F
  - determination of c_B

Outlook:

- heavy quark mass region
- systematic errors controlled as $a \to 0$
- efficient procedures for full QCD
- matching with isotropic lattice
- applications to various heavy-light matrix elements.