Anisotropic lattice with nonperturbative accuracy

Hideo Matsufuru

High Energy Accelerator Research Organization (KEK), Japan

Hidenori Fukaya, Tetsuya Onogi, Takashi Umeda

(Yukawa Institute for Theoretical Physics, Kyoto Univ.)

Masanori Okawa

(Dept. Physics, Hiroshima Univ.)

Web-site: http://www.rcnp.osaka-u.ac.jp/~matufuru/

Lattice 2004

21–26 June 2004, FNAL, US

Contents

Anisotropic lattice for heavy quark physics

- Introduction
  — motivation, our approach, present status
- Calibration of gauge field
- Calibration of quark field
- Summary and outlook

Introduction

Our motivation:
high precision computation of hadronic matrix elements for flavor physics

Recent experimental development of in heavy flavor physics such as B factories, Charm factory
$\longrightarrow$ need precise theoretical prediction to a few percent level

Matrix elements:

- decay constants
- bag parameters
- form factors
- quark masses
- light-cone wave functions

etc.

Our approach: why anisotropic lattice?

Problem in lattice QCD calculations:
for heavy quarks (c and b), large $O(am_Q)$ error

For precision computation of heavy-light matrix elements,
we need a framework for heavy quark which has

(i) continuum limit
— to remove lattice artifact

(ii) systematic improvement
such as nonperturbative renormalization technique
(for $m_q \simeq 0$)

(iii) modest size of computation

Our approach: why anisotropic lattice?

Previous approaches satisfy not all of them:

a). Effective theories (NRQCD, etc) — (i)$\times$

Thacker and Lepage, Phys. Rev. D 43 (1991) 196.

b). Relativistic framework — (iii)$\times$

c). Fermilab approach (with/without mass dependent tuning) — (ii)$\times$

El-Khadra et al., Phys. Rev. D 55 (1997) 3933.

Sroczynski, et al., Nucl. Phys. B (PS) 83 (2000) 971.

So far, systematic uncertainty $\sim$ 10%

To achieve calculations with $\sim$2% precision, we need yet another approach which satisfies above conditions (i)–(iii).

$\Rightarrow$ our proposal: Anisotropic lattice

Our approach

Anisotropic lattice: $a_\tau < a_\sigma \Rightarrow$ Anisotropy $\xi = a_\sigma / a_\tau$

Quark action: $O(a)$ improved, along with Fermilab approach

- □ bare anisotropy parameter γ_F:
  in general, to be tuned mass dependently
  e.g., using meson dispersion relation

- ○ Continuum limit – *OK*
- ○ Modest computational cost – *OK*
- ○ Systematic improvement ?
  If quark mass is sufficiently less than a^{-1},
  tuned parameters for massless quark are applicable.
  $\Rightarrow$ nonperturbative renormalization technique
  Lüscher et al., Nucl.Phys. B 491 (1997) 323.

Our approach

Our expectation:

For $m_Q \ll a_\tau^{-1}$, (not necessarily $m_Q \ll a_\sigma^{-1}$),
mass dependences of parameters in the action are so small that the tuned parameters for massless quark are also applicable in such a quark mass region.

— Then, systematic improvement is possible.
(performed at $m_q \simeq 0$)

$\Rightarrow$ To be justified numerically, and in perturbation theory.

- Tree level: *OK*
- $O(a^0)$ improved version: *OK* (2% accuracy)
- High precision: in progress

Present status

□ One-loop perturbative calculation:

Harada et al., Phys. Rev. D 64 (2001) 074501

Renormalization factors of heavy-light bilinears and quark rest mass at $m_Q a_\sigma \sim 1$, $m_Q \ll a_\tau^{-1}$

$\Rightarrow$ Well approximated with linear form in $m_Q a_\tau$

— quark mass dependence can be controlled

□ Numerical simulation:

In quenched approximation, tadpole improved c_E and c_B

- Mass dependent tuning

Matsufuru, Onogi and Umeda, Phys. Rev. D 64 (2001) 114503

γ_F is tuned with meson dispersion relation

Quark mass dependence is small for $m_q a_\tau \ll 1$

Present status (cont.)

- Test of relativity relation

  Harada et al., Phys. Rev. D 66 (2002) 014509

  Heavy-light meson dispersion relation for $\gamma_F = \gamma_F(m_q = 0)$
  Relativity relation well holds for $m_q a_\tau \ll 1$ (while $m_q a_\sigma \sim 1$)

- Application to decay constant

  Matsufuru, Harada, Onogi and Sugita, hep-lat/0209090

  – Around charm quark mass, with $O(10\%)$ accuracy
  – f_{D_s} consistent with precision computation
  by ALPHA Collab.
  – β dependence of f_{D_s}/f_K very small.
  – Result also consistent with previous works

— Encouraging results for further development

Toward high precision computations

Calibrations of parameters to 0.2% level of accuracy
in quenched approximation

$\Rightarrow$ Applicable to a few percent calculation of matrix elements

□ Gauge field:

- Precise computation of static quark potential
- Renormalized anisotropy defined with r_0

$\Rightarrow$ $O(0.2\%)$ calibration is possible

□ Quark field:

γ_F, c_E, c_B, and c_A (for axial current) in massless limit.

— γ_F must be tuned precisely (to $O(0.2\%)$ level)

Combine two procedures:

- Nonperturbative renormalization technique
- Spectroscopy in fine and coarse directions

— Numerical simulation is in progress

Calibration of gauge field

Previous result for gauge field:
Klassen's work with 1% level of statistical error.

T.R.Klassen, Nucl. Phys. B 533 (1998) 557.

— For present purpose, this accuracy is not sufficient.

Renormalized anisotropy $\xi_G \leftarrow$ static potential
We define ξ_G through hadronic radius r_0

r_0: $r_0^2 F(r_0) = 1.65$ ($F(r)$: force)

Sommer, Nucl. Phys. B411 (1994) 839

- Precise computation possible
- Good scaling behavior (if lattice scale is set by r_0)

Calibration scale is unambiguous.
Continuum limit in terms of scale set by r_0
$\rightarrow$ systematic error in γ_G disappers as $a \rightarrow 0$

Calibration of gauge field

Precise computation of static potential:

- Lüscher-Weisz noise reduction technique

  Lüscher and Weisz, JHEP 0109 (2001) 010.

  – accurate computation is possible.
  – rather large memory is required.
- Smearing technique in anisotropic plane
  – standard technique on isotropic lattices
  – convenient for large lattices

Target anisotropy: $\xi = 4$

$\beta = 5.7 - 6.3$ $(a_s^{-1} = 1 - 3$ GeV)

At each β, several input $\gamma_G \rightarrow \gamma_G^*$ s.t. $\xi_G(\gamma_G^*) = \xi$

where $\xi_G = r_0^{(fine)}/r_0^{(coarse)}$

Calibration of gauge field

Result at $\beta = 5.75$





$\gamma^* = 3.1396(76)$ (0.24% accuracy)
— almost sufficient (further improvement is easy)

Calibration of gauge field

Result at $\beta = 6.00$





$\gamma^* = 3.227(25)$ (0.8% accuracy)

— not sufficient, can be improved by statistics

Calibration of gauge field

Results (preliminary)



More accuracy is needed in wide range of β (work in progress).

global fit $\rightarrow$ γ_G^* with 0.2% level uncertainty

Calibration of quark field

Quark action: $O(a)$ improved Wilson action

$$S_F = \sum_{x,y} \bar{\psi}(x) K(x,y) \psi(y)$$

$$\begin{aligned} K(x,y) &= \delta_{x,y} - \kappa_\tau \left[(1-\gamma_4) U_4(x) \delta_{x+\hat{4},y} + (1+\gamma_4) U_4^\dagger(x-\hat{4}) \delta_{x-\hat{4},y} \right] \\ &\quad - \kappa_\sigma \sum_i \left[(r-\gamma_i) U_i(x) \delta_{x+\hat{i},y} + (r+\gamma_i) U_i^\dagger(x-\hat{i}) \delta_{x-\hat{i},y} \right] \\ &\quad - \kappa_\sigma c_E \sum_i \sigma_{4i} F_{4i}(x) \delta_{x,y} + r \kappa_\sigma c_B \sum_{i>j} \sigma_{ij} F_{ij}(x) \delta_{x,y} \end{aligned}$$

$\gamma_F = \kappa_\tau / \kappa_\sigma$: bare anisotropy

γ_F, c_E and c_B: to be tuned nonperturbatively

Calibration of quark field

Five parameters (γ_F, c_E, c_B, c_A, κ_c) should be determined.

Calibration steps:

(1) Schrödinger functional method
$\Rightarrow$ tuning of c_E, c_B

(2) Spectrum in coarse / fine directions on lattices of T,$L \sim 2$ fm
Physical isotropy conditions for m_{PS}, m_V $\Rightarrow$ γ_F, (c_B)

(3) Schrödinger functional method
$\Rightarrow$ determine c_A, κ_c

(4) Check of systematic errors
Light hadron spectrum and dispersion relation
Taking the continuum limit

Calibration of quark field

Determination of c_E:

Nonperturbative improvement technique

Lüscher et al., Nucl. Phys. B478 (1996) 365

Lüscher et al., Nucl. Phys. B491 (1997) 323

Schrödinger functional method:
implementation for anisotropic lattice is straightforward.
(t-direction set to the fine direction)

PCAC relation up to $O(a^2) \rightarrow$ improvement of c_E

Unrenormalized quark mass
$m = \frac{1}{2}[\frac{1}{2}(\partial_0^* + \partial_0) f_A(t) + c_A a_t \partial_0^* \partial_0 f_P(t)]/f_P(t)$

Improvement condition:
m does not depend on kinematical parameters such as boundary gauge field.

Calibration of quark field

Result at high β



$\beta = 9.5$, $8^3 \times 64$, $\xi = 4$ (gluonic anisotropy from Klassen's work)

- NP improved c_E is close to 1-loop mean-field value, c_E=1.13.
- ΔM is almost independent of c_B

Calibration of quark field

Result at $\beta = 6.10$



$\beta = 6.10$, $8^3 \times 64$, $\xi = 4$ ($a_s^{-1} = 2.0$ GeV) (γ_G by Klassen)

- NP improved c_E is larger than tadpole tree value, $c_E = 1.26$.
- ΔM is not sensitive to c_B

Calibration of quark field

Determination of γ_F:



$\beta = 5.75$, $12^2 \times 24 \times 96$, $a_s^{-1} = 1.1$ GeV, 136 confs.

γ_F is determined from meson masses in fine/coarse directions

- precision of γ_F is still not sufficient
- consistent with γ_F from dispersion relation

Calibration of quark field

Determination of c_B:



In principle, c_B can be determined by mass splitting in fine/coarse directions.

In practice, signal-to-noise ratio is too large.

We need other procedures:

- Schrödinger functional with boundaries in coarse direction

Summary

We are developing anisotropic lattices for precise computation of heavy-light matrix elements.

Results so far are encouraging for further development.

Now calibration of gauge and quark fields are in progress:

- high precision calibration for gluonic anisotropy
- nonperturbative determination of c_E

while we still need improved procedures for

- accurate determination of γ_F
- determination of c_B

Outlook

We also need to demonstrate:

- the parameters at $m_q = 0$ suffices for heavy quarks
- all systematic errors can be controlled as $a \to 0$

and to develop

- efficient procedures for full QCD
- matching with isotropic lattice

in addition to application to various heavy-light matrix elements.