

**AAC ANALYSIS OF POLARIZED PARTON DISTRIBUTIONS
WITH UNCERTAINTIES**

M. HIRAI*, S. KUMANO†, N. SAITO‡
(ASYMMETRY ANALYSIS COLLABORATION)

* *Institute of Particle and Nuclear Studies, KEK
1-1, Oho, Tsukuba, 305-0801, Japan*

† *Department of Physics, Saga University, Saga, 840-8502, Japan*

‡ *Department of Physics, Kyoto University, Kyoto, 606-8502, Japan*

We report recent studies of the Asymmetry Analysis Collaboration (AAC) on polarized parton distribution functions (PDFs). Using the data on the spin symmetry A_1 in deep inelastic lepton scattering, we investigate optimum polarized PDFs. Their uncertainties are estimated by the Hessian method. The uncertainties are large for the polarized antiquark and gluon distributions. We discuss the role of accurate SLAC-E155 proton data on the determination of the PDFs. The obtained distributions are compared with other parametrization results.

1 Introduction

Polarized parton distribution functions (PDFs) have been investigated by using experimental data on the spin asymmetry A_1 in deep inelastic lepton-nucleon scattering. Thanks to several experiments, gross properties of the polarized PDFs are now known. However, the nucleon spin structure is still far from complete understanding. It becomes clear that the fraction of nucleon spin carried by quarks is small; however, the gluon contribution cannot be specified at this stage. Although the nucleon spin is one of fundamental physics quantities, we do not know how the spin consists of quark and gluon spins.

Because the polarized distributions cannot be completely fixed at this stage, it is especially important to show their uncertainties for summarizing the present knowledge and for planning future experimental projects. Such uncertainties have been investigated in the unpolarized PDFs in the last several years, and there were also recent error analyses for the polarized ones. We report a new analysis by the Asymmetry Analysis Collaboration (AAC) for the polarized PDFs and their uncertainties [1].

The AAC proposed the first parametrization in Ref. [2]. The additional data to the first AAC analysis (AAC00) are the SLAC-E155 proton data. Because they are accurate, it is interesting to investigate their effects on the determination of the polarized distributions. Namely, we discuss how the data reduce the uncertainties of the polarized PDFs. Next, the global analysis was repeated by taking $\Delta g=0$ at $Q^2=1 \text{ GeV}^2$ in order to show error correlation effects between the polarized antiquark and gluon distributions.

This paper consists of the following. In Sec. 2, a method is discussed for obtaining the polarized PDFs and their uncertainties by analyzing A_1 data. Analysis results are shown in Sec. 3, and they are summarized in Sec. 4.

2 Analysis method

The polarized PDFs are determined by analyzing the data of spin asymmetry for inclusive lepton scattering in the deep inelastic region. The spin asymmetry A_1 is expressed by the structure functions, F_2 and g_1 , and the longitudinal-transverse ratio R :

$$A_1(x, Q^2) = \frac{g_1(x, Q^2)}{F_2(x, Q^2)} 2x [1 + R(x, Q^2)]. \quad (1)$$

The function F_2 is expressed in terms of unpolarized PDFs, for which we use the GRV98 parametrization. For the function R , a SLAC analysis result is used. The function g_1 is expressed in terms of the polarized distributions:

$$g_1(x, Q^2) = \frac{1}{2} \sum_{i=1}^{n_f} e_i^2 \left\{ \Delta C_q(x, \alpha_s) \otimes [\Delta q_i(x, Q^2) + \Delta \bar{q}_i(x, Q^2)] \right. \\ \left. + \Delta C_g(x, \alpha_s) \otimes \Delta g(x, Q^2) \right\}, \quad (2)$$

where the symbol \otimes indicates the convolution integral $f(x) \otimes g(x) = \int_x^1 dy f(x/y)g(y)/y$. The functions Δq , $\Delta \bar{q}$, and Δg are polarized quark, antiquark, and gluon distributions. The ΔC_q and ΔC_g are polarized coefficient functions in the next-to-leading order.

The functional form of the polarized PDFs is chosen at the initial Q^2 ($\equiv Q_0^2$):

$$\Delta f(x) = [\delta x^\nu - \kappa(x^\nu - x^\mu)]f(x), \quad (3)$$

where $f(x)$ is the unpolarized PDF. The factors δ , ν , κ , and μ are the parameters to be determined by the χ^2 analysis. For Δf , we take Δu_v , Δd_v , $\Delta \bar{q}$, and Δg by assuming flavor symmetric antiquark distributions ($\Delta \bar{u} = \Delta \bar{d} = \Delta \bar{s} \equiv \Delta \bar{q}$) at Q_0^2 . The initial scale is taken $Q_0^2=1 \text{ GeV}^2$. The distributions are evolved to the experimental Q^2 points so that the following χ^2 is calculated:

$$\chi^2 = \sum_i \frac{[A_{1,i}^{\text{data}}(x, Q^2) - A_{1,i}^{\text{theo}}(x, Q^2)]^2}{[\Delta A_{1,i}^{\text{data}}(x, Q^2)]^2}, \quad (4)$$

where $A_{1,i}^{\text{data}}$ and $A_{1,i}^{\text{theo}}$ indicate an asymmetry data and a calculated asymmetry, and $\Delta A_{1,i}^{\text{data}}$ is an experimental error. The actual χ^2 minimization is done by the CERN program library, MINUIT. The data are taken by the EMC, SMC, SLAC-E130, E142, E143, E154, E155, and HERMES collaborations. Recent JLab data will be included in our future analysis.

Uncertainties of the polarized PDFs are calculated by the Hessian method. Running the MINUIT subroutine, we obtain the Hessian matrix H , which is then used for calculating the uncertainties:

$$[\delta F(x)]^2 = \Delta \chi^2 \sum_{i,j} \left(\frac{\partial F(x, a)}{\partial a_i} \right)_{a=\hat{a}} H_{ij}^{-1} \left(\frac{\partial F(x, a)}{\partial a_j} \right)_{a=\hat{a}}. \quad (5)$$

Here, F indicates a polarized PDF, a_i is a parameter in the χ^2 fit, and \hat{a} indicates the optimized parameter set. The $\Delta \chi^2$ value determines the confidence level, and it is chosen as $\Delta \chi^2=12.647$ by considering that the number of parameters is eleven [1]. It corresponds to the one- σ -error range.

3 Results

The optimum polarized PDFs obtained by the χ^2 fit are shown in Fig. 1. The uncertainties are shown by the shaded areas for the AAC03 version and by the dashed curves for the previous version AAC00. The difference between AAC00 and AAC03 is the addition of the accurate E155 proton data. It is obvious from the figure that the accurate E155 data improves the PDF determination. The magnitudes of the uncertainty bands are significantly reduced. It is noteworthy that the gluon uncertainties are also reduced in spite of the fact that the polarized gluon contribution to g_1 is rather small. It should be caused by the error correlation between the antiquark and gluon distributions. In order to investigate such a correlation, we repeated the global analysis by terminating the gluon distribution ($\Delta g=0$) at the initial Q^2 (Q_0^2). Then, we find that the uncertainties of $\Delta\bar{q}$ are significantly reduced, while those of Δu_v and Δd_v remain unchanged. It suggests that the errors of the polarized antiquark and gluon distributions should be strongly correlated. From these studies, it is clear that an accurate determination of Δg is important for the determination of $\Delta\bar{q}$ and vice versa.




Figure 1. Polarized PDFs and their uncertainties are shown at $Q^2=1 \text{ GeV}^2$. The solid curves and shaded areas indicate the AAC03 distributions and uncertainties. The dashed curves indicate the uncertainties of the previous version AAC00.

The AAC03 distributions are compared with some of other analysis results, GRSV [3], LSS [4], and BB [5], in Fig. 2. All the distributions roughly agree, and they are within the uncertainty bands. The uncertainties of Δg are especially large, which indicates that even $\Delta g = 0$ is allowed at this stage. There are differences between the antiquark distributions at small x , and it leads to different values of quark spin content [2].




Figure 2. The polarized PDFs are compared with other parametrizations at $Q^2=1 \text{ GeV}^2$.

4 Summary

We reported the AAC03 analysis for determining polarized parton distribution functions. The uncertainties were estimated by the Hessian method. Comparing the AAC03 results with the previous ones (AAC00), we find that the E155 proton data played an important role in reducing not only the uncertainties of Δq and $\Delta \bar{q}$ but also the Δg uncertainties due to the error correlation between $\Delta \bar{q}$ and Δg . The uncertainties of Δg are huge, so that we need future experimental measurements such as prompt-photon measurements at RHIC [6].

Acknowledgements

S.K. was supported by the Grant-in-Aid for Scientific Research from the Japanese Ministry of Education, Culture, Sports, Science, and Technology.

References

1. M. Hirai, S. Kumano, and N. Saito (Asymmetry Analysis Collaboration), Phys. Rev. D69 (2004) 054021. URL: <http://spin.riken.bnl.gov/aac/>.
2. Y. Goto *et al.* (AAC), Phys. Rev. D62 (2000) 034017.
3. M. Glück, E. Reya, M. Stratmann, and W. Vogelsang, Phys. Rev. D63 (2001) 094005.
4. E. Leader, A. V. Sidorov, and D. B. Stamenov, Eur. Phys. J. C23 (2002) 479; A. V. Sidorov, contribution to this workshop.
5. J. Blümlein and H. Böttcher, Nucl. Phys. B636 (2002) 225.
6. M. Hirai, hep-ph/0404279 (Phys. Lett. B in press).